

BAFF/TNFSF13B Protein, Human, Recombinant (CHO)

General Information

Synonyms:	BLYS;B-cell Activating Factor;TALL-1;THANK;TALL1;CD257;TNFSF13B;TNFSF20;ZTNF4
Protein Construction:	Ala134-Leu285
Species:	Human
Expression Host:	CHO Cells
Accession:	Q9Y275
Molecular Weight:	~17 kDa (Non-reducing conditions)

QC Testing

Biological Activity:	ED 50 < 20.0 ng/ml, determined by dose-dependent mitogenic activity on human RPMI 8226 cells, corresponding to a specific activity of > 5.0 x 10 ⁴ units/mg.
Purity:	> 95% as determined by SDS-PAGE; > 95% as determined by HPLC
Endotoxin:	< 0.2 EU/μg of protein as determined by the LAL method.
Formulation:	Lyophilized from a 0.2 μm filtered solution in PBS.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

B-cell activating factor, also known as BAFF, TALL-1, TNFAK, and zTNF4, is a member of the TNF ligand superfamily designated TNFSF13B. Produced by macrophages, dendritic cells, and T lymphocytes, BAFF promotes the survival of B cells and is essential for B cell maturation. BAFF binds to three TNF receptor superfamily members: B-cell maturation antigen (BCMA/TNFRSF17), transmembrane activator and calcium-modulator and cyclophilin ligand interactor (TACI/TNFRSF13B) and BAFF receptor (BAFF R/BR3/TNFRSF 13C). These receptors are type III transmembrane proteins lacking a signal peptide. Whereas TACI and BCMA bind BAFF and another TNF superfamily ligand, APRIL (a proliferation-inducing ligand), BAFF R selectively binds BAFF. The BAFF R extracellular

A DRUG SCREENING EXPERT

domain lacks the TNF receptor canonical cysteine-rich domain (CRD) and contains only a partial CRD with four cysteine residues. Human and mouse BAFF R share 56% aa sequence identity. BAFF R is highly expressed in spleen, lymph node and resting B cells. It is also expressed at lower levels in activated B cell, resting CD4 + T cells, thymus and peripheral blood leukocytes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481