

Anti-Flag Tag Antibody (2H702)

Product Details

Ig Type:	IgG1
Reactivity:	Species independent
Molecular Weight:	Actual: 48 kDa.
Clone:	2H702
Purification:	Protein G purified

Applications

1. Sample:

DDDDK-Tagged Fusion Protein Overexpression E.coli Lysate

Primary: Anti-Flag Tag (TMAB-00684) at 1/5000 ~ 1/100000 dilution



Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution

Predicted band size: 41 kDa

Observed band size: 45 kDa

Verified Activity:

2. 20 ng Recombinant human TNFSF13B protein (HEK293), N-hFc-Flag per Lane probed with Flag Tag monoclonal antibody respectively, unconjugated (TMAB-00684) at 1:50000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at RT for 60 min.



Application: ELISA,FCM,ICC/IF,IF,IHC-Fr,IHC-P,WB

Recommended

ELISA=1:5000-10000; FCM=1 µg/Test; ICC/IF=1:100-500; IF=1:200-1000; IHC-Fr=1:200-1000; IHC-P=1:200-1000; WB=1:10000-100000

A DRUG SCREENING EXPERT

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated Flag Tag

Synonyms: Enterokinase Cleavage Site tag;DYKDDDDK epitope tag;DYKDDDDK tag;FLAG;ECS epitope tag; ECS tag;DDDK;DYKDDDDK;Enterokinase Cleavage Site epitope tag;ddk;DDDDK epitope tag

Biology Area: DDDDK Tag

Research Background

This is a useful tool for the localisation and characterisation of DDDDK tagged proteins.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481