

Anti-Phospho-NFKB1 (Ser337) Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human, Mouse (predicted: Rat, Pig, Cow)
Molecular Weight:	Theoretical: 48/105 kDa. Actual: 48 kDa.
Purification:	Protein A purified

Applications

1. Sample:

Raji (Human) Cell Lysate at 30 µg

Primary: Anti-Phospho-NFKB1 (Ser337) (TMAB-01470) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 48/105 kDa

Observed band size: 48 kDa

2. Blank control (blue line): A549 (fixed with 70% ethanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice).

Primary Antibody (green line): Rabbit Anti-Phospho-NFKB1 (Ser337) antibody (TMAB-01470), Dilution: 1 µg/10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG.

Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC, Dilution: 1 µg/test.

3. Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 min; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-NFKB1 (Ser337)) Polyclonal Antibody, Unconjugated (TMAB-01470) at 1:200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

Verified Activity:

4. Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 min; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-NFKB1 (Ser337)) Polyclonal Antibody, Unconjugated (TMAB-01470) at 1:200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

5. Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 min; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-NFKB1 (Ser337)) Polyclonal Antibody, Unconjugated (TMAB-01470) at 1:400 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

6. Sample:

U87-MG (Human) Lysate at 30 µg







MCF7 (Human) Lysate at 30 µg

Primary: Anti-Phospho-NFKB1 (Ser337) (TMAB-01470) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 48/105 kDa

Observed band size: 48/105 kDa



Application: FCM,IF,IHC-Fr,IHC-P,WB

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1µg/Test

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated Synthesised phosphopeptide: human NF KappaB p105 around the phosphorylation site of Ser337

Antigen Species: Human

Gene ID: 4790

Uniprot ID: P19838

Synonyms: p-NFKB1 (S337);Nuclear factor NF-kappa-B p105 subunit;p-NFKB1 (Ser337);NFKB1;EBP-1;NFKB1 (p-S337);NFKB1 (p-Ser337);Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1;DNA-binding factor KBF1

Biology Area: p50,NFKB pathway,p50,NFkB pathway,Transcription Factors,Macrophage / Inflamm.,NFkB Pathway

Research Background

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481