

Anti-SLC17A6 Polyclonal Antibody



Product Details

| | |
|-------------------|---|
| Ig Type: | IgG |
| Reactivity: | Rat (predicted:Human,Mouse,Chicken,Dog,Pig,Cow,Horse) |
| Molecular Weight: | Theoretical: 64 kDa. |
| Purification: | Protein A purified |

Applications

1. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-VGLUT2 Polyclonal Antibody, Unconjugated (TMAB-01704) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining.
2. Positive sample: Paraformaldehyde-fixed, paraffin embedded (rat spinal cord); Negative sample: Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 min; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (VGLUT2) Polyclonal Antibody, Unconjugated (TMAB-01704) at 1:200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

Verified Activity:



| | |
|--------------|--|
| Application: | IF,IHC-Fr,IHC-P |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated synthetic peptide: human VGLUT2

Antigen Species: Human

Gene ID: 57084

Uniprot ID: Q9P2U8

Synonyms: VGLUT2;Solute carrier family 17 member 6;Differentiation-associated BNPI;Differentiation-associated Na(+)-dependent inorganic phosphate cotransporter;Vesicular glutamate transporter 2;Slc17a6;Dnpi

Biology Area: Transporters

Research Background

The ATP-dependent, chloride-sensitive vesicular glutamate transporters (VGLUT) include BNPI (VGLUT1), VGLUT2 (DNPI) and VGLUT3. The brain expresses BNPI (brain specific Na⁺-dependent inorganic phosphate (Pi) cotransporter) and VGLUT2 in a complementary fashion. The telencephalic regions express BNPI, whereas the lower brainstem and diencephalic regions express VGLUT2. Rat pinealocytes express both BNPI and VGLUT2. The striatum, hippocampus, cerebral cortex and raphe nuclei express VGLUT3 in a small number of neurons. Pancreatic a and b cells express BNPI and VGLUT2 in response to glucose concentrations. Human VGLUT3 shares a 72% sequence homology with VGLUT2 and BNPI.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481