

Anti-CDK2 Antibody (3L593)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 33 kDa.
Clone:	3L593
Purification:	ProA affinity purified

Applications

- Verified Activity:
1. Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Cdk2 antibody. Counter stained with hematoxylin.
 2. Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-Cdk2 antibody. Counter stained with hematoxylin.
 3. Immunohistochemical analysis of paraffin-embedded mouse spleen tissue using anti-Cdk2 antibody. Counter stained with hematoxylin.
 4. ICC staining Cdk2 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
 5. ICC staining Cdk2 in CRC cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
 6. ICC staining Cdk2 in SH-SY-5Y cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
 7. ICC staining Cdk2 in N2A cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
 8. All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.
 9. All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.
 10. All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.





