

Anti-Phospho-HER2/ERBB2 (Thr 686) Antibody (1M462)

Product Details

Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 185 kDa.
Clone:	1M462
Purification:	ProA affinity purified

Applications

Verified Activity: 1. Western blot analysis of Neu phosphorylation in serum starved A431 (A), and serum starved A431 treated for 15 minutes with PMA (B), LPA (C), Ceramide (D), Bradykinin (E) and Bombesin (F) whole cell lysates.



Application:	IP,WB
Recommended	WB: 1:100-1000; IP: 1-2 µg per 100-500 µg of total protein(1 ml of cell lysate)

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	Amino acid residues surrounding Threonine 686 of Neu of human origin
Antigen Species:	Human
Uniprot ID:	P04626
Synonyms:	p-HER2/ERBB2 (T 686);HER2/ERBB2 (p-T 686);HER2/ERBB2 (p-Thr 686);p-HER2/ERBB2 (Thr 686)

Research Background

Neu (ErbB-2 erythroblastic leukemia viral oncogene homolog 2, HER-2, NGL, TKR1, c-erb B2) oncogene was originally cloned from a rat neuroglioblastoma. Human Neu is referred to as HER-2 since the protein structure resembles human epidermal growth factor receptor (HER). ErbB-2 refers to a high level of similarity to ErbB (avian erythroblastosis oncogene B), later found to code for EGFR (HER). Tyr 1248-phosphorylated Neu localizes with Mucin 4/sialomucin complex at the apical surfaces of ductal and alveolar cells in rodent lactating gland. Phosphorylation of Neu at Tyr 1139 promotes association of GRB2 and GRB7 through an Src homology 2 (SH2) domain-dependent interaction and contributes to the etiology of certain breast, gastric and esophageal cancers and testicular germ cell

tumors. Neu phosphorylation on Tyr 1221 and Tyr 1248 promotes association of Shc (SH2 domain-containing transforming protein 1) through an SH2 domain. Neu phosphorylation at Tyr 1196 and Tyr 1248 promotes association of Shc through a PTB (phosphotyrosine binding) domain. SH2 and PTB domains recognize tyrosine phosphorylated proteins in a sequence-specific fashion and transduce extracellular signals via subcellular targeting, directing assembly of complexes and modulating enzymatic activity.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481