

Anti-MAP2K3/MAP2K6 Antibody (7F635)



Product Details



Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 39/37 kDa.
Clone:	7F635
Purification:	ProA affinity purified

Applications

Verified Activity:

1. Western blot analysis of MEK3+MEK6 on different lysates using anti-MEK3+MEK6 antibody at 1/1,000 dilution. Positive control: Lane 1: Hela, Lane 2: Jurkat.
2. Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-MEK3+MEK6 antibody. Counter stained with hematoxylin.
3. ICC staining MEK3+MEK6 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
4. ICC staining MEK3+MEK6 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
5. ICC staining MEK3+MEK6 in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
6. Flow cytometric analysis of Hela cells with MEK3+MEK6 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.





Application: FCM, ICC/IF, IHC, IP, WB

Recommended WB: 1:1000-2000; IHC: 1:50-200; ICC/IF: 1:50-200; FCM: 1:50-100

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein

Uniprot ID: P46734

Synonyms: PRKMK6; ERK kinase 6; MEK6; MAPKK 6; MKK6; MPK 6; Mitogen activated protein kinase kinase 3; MAP2K 6; zMKK 3; MAPK kinase 6; MAPK ERK kinase 3; MAPK ERK kinase 6; MKK3; mMKK 3b; MAP kinase kinase 3; SKK2; PRKMK 3; MEK 3; Protein kinase mitogen activated kinase 6; MEK3; MAPKK3; MAP Kinase Kinase 6; Mitogen activated protein kinase kinase 6; mMKK3b; mMKK 6b; ERK kinase 3; MAPKK 3; MAPK kinase 3; MAPKK6; Dual specificity mitogen activated protein kinase kinase 3;

PRKMK 6;Dual specificity mitogen activated protein kinase kinase 6;Protein kinase mitogen activated kinase 3;PRKMK3;zMKK 6;MEK 6;MKK 6;MAP2K 3;MKK 3;MP2K6;MP2K3;MPK 3; mMKK6b

Research Background

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481