

Anti-MYD88 Antibody (2C378)



Product Details





Ig Type:	IgG
Reactivity:	Human
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 33 kDa.
Clone:	2C378
Purification:	ProA affinity purified

Applications

Verified Activity:

1. Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-MyD88 antibody. Counter stained with hematoxylin.
2. Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-MyD88 antibody. Counter stained with hematoxylin.
3. ICC staining MyD88 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
4. ICC staining MyD88 in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
5. ICC staining MyD88 in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.
6. Flow cytometric analysis of Hela cells with MyD88 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.





Application: FCM, ICC/IF, IHC, WB

Recommended WB: 1:1000; IHC: 1:50-200; ICC/IF: 1:100-500; FCM: 1:50-100

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein

Uniprot ID: Q99836

Synonyms: MYD88; Myeloid differentiation primary response protein MyD88

Research Background

Interleukin-1 (IL-1)-induced activation of the NFκB pathway is mediated through the IL-1 receptor and the subsequent phosphorylation of IL-1 receptor-associated kinase (IRAK). The myeloid differentiation protein MyD88 was originally characterized as a protein upregulated in myeloleukemic cells following IL-6-induced growth arrest and terminal differentiation. MyD88 is now known to function as an adaptor protein for the association of IRAK with the IL-1 receptor. MyD88 is functionally homologous to the adaptor protein tube in the Toll signaling pathway of

A DRUG SCREENING EXPERT

Drosophila, and both proteins are members of the Toll/IL-1R superfamily. MyD88 contains a characteristic N-terminal death domain that is essential for NFκB activation and an adjacent Toll/IL-1R homology domain (TIR domain). Collectively, these domains enable the protein-protein interactions of MyD88 with IRAK and the IL-1 receptor complex.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481