

Anti-NDUFS4 Antibody (3X649)




Product Details


Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 20 kDa.
Clone:	3X649
Purification:	ProA affinity purified

Applications

Verified Activity:

1. Western blot analysis of Ndufs4 on rat heart tissue lysate using anti-Ndufs4 antibody at 1/1,000 dilution.
2. Immunohistochemical analysis of paraffin-embedded rat kidney tissue using anti-Ndufs4 antibody. Counter stained with hematoxylin.
3. Immunohistochemical analysis of paraffin-embedded human small intestine tissue using anti-Ndufs4 antibody. Counter stained with hematoxylin.
4. Flow cytometric analysis of SH-SY-5Y cells with Ndufs4 antibody at 1/100 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.





Application: FCM,IHC,IP,WB

Recommended WB: 1:500-2000; IHC: 1:50-200; IP: 1:10-50; FCM: 1:50-100

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: human Ndufs4 aa 43-175

Antigen Species: human

Uniprot ID: O43181

Synonyms: mitochondrial;NADH-ubiquinone oxidoreductase 18 kDa subunit;NDUFS 4;Complex I-AQDQ;CI-AQDQ;CI-18 kDa;Complex I-18 kDa;NADH dehydrogenase [ubiquinone] iron-sulfur protein 4

Research Background

Complex 1 (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex 1 is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFS4 (NADH dehydrogenase (ubiquinone) Fe-S protein 4), also known as AQDQ or CI-18 (Complex I-18kDa protein), belongs to the Complex I NDUFS4 subunit family. NDUFS4 localizes to the matrix side of the inner membrane of the mitochondrion and functions as an accessory subunit of Complex I. Mutations in the gene encoding NDUFS4 can result in Complex I mitochondrial respiratory chain deficiency. Patients with this deficiency may exhibit cardiomyopathy, myopathy, liver failure and neurological disorders.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481