

## Anti-Phospho-NFATc4 (Ser168, 170) Polyclonal Antibody

### Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse
Conjugation:	Unconjugated
Molecular Weight:	Actual: 140 kDa.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

### Applications

Application:	ELISA,IF,IHC-P,WB
Recommended	WB: 1:500-2000; IHC-P: 1:100-300; ELISA: 1:5000

### Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen:	A synthesized phosphopeptide: human NFAT3 around the phosphorylation site of Ser168 and Ser170. AA range:136-185
Antigen Species:	Human
Uniprot ID:	Q14934
Synonyms:	NFATc4 (p-S168, 170);p-NFATc4 (S168, 170);p-NFATc4 (Ser168, 170);NFATc4 (p-Ser168, 170)

### Research Background

nuclear factor of activated T-cells 4(NFATC4) Homo sapiens This gene encodes a member of the nuclear factor of activated T cells (NFAT) protein family. The encoded protein is part of a DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor stimulation and an inducible nuclear component. NFAT proteins are activated by the calmodulin-dependent phosphatase, calcineurin. The encoded protein plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of interleukin-2 and interleukin-4. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014],

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481