

Anti-Phospho-GRIN1 (Ser890) Polyclonal Antibody 2



Product Details

| | |
|-------------------|--|
| Ig Type: | IgG |
| Reactivity: | Human,Mouse,Rat |
| Conjugation: | Unconjugated |
| Molecular Weight: | Actual: 120 kDa. |
| Purification: | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide. |

Applications

Verified Activity:

1. Immunohistochemical analysis of paraffin-embedded human brain tissue using NMDAR1 (Phospho-Ser890) antibody TMAC-02857 (left) or the same antibody preincubated with blocking peptide (right).
2. Immunofluorescence staining of methanol-fixed A549 cells using NMDAR1 (Phospho-Ser890) Antibody TMAC-02857.



| | |
|--------------|------------------------------|
| Application: | IF,IHC |
| Recommended | IHC: 1:50-100; IF: 1:100-200 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | Peptide sequence around phosphorylation site of Serine 890(A-S-S(p)-F-K) derived from Human NMDAR1 |
| Antigen Species: | Human |
| Uniprot ID: | Q05586 |
| Synonyms: | p-GRIN1 (S890);GRIN1 (p-Ser890);p-GRIN1 (Ser890);GRIN1 (p-S890) |

Research Background

NMDA receptors are members of the ionotropic class of glutamate receptors, which also includes Kainate and AMPA receptors. NMDA receptors consist of NR1 subunits combined with one or more NR2 (A-D) or NR3 (A-B) subunits. The ligand-gated channel is permeable to cations including Ca^{2+} , and at resting membrane potentials NMDA receptors are inactive due to a voltage-dependent blockade of the channel pore by Mg^{2+} . NMDA receptor activation, which requires binding of glutamate and glycine, leads to an influx of Ca^{2+} into the postsynaptic region where it activates several signaling cascades, including pathways leading to the induction of long-term potentiation (LTP) and depression (LTD). NMDA receptors have a critical role in excitatory synaptic transmission and plasticity in the CNS. They govern a range of physiological conditions including neurological disorders caused by excitotoxic neuronal injury, psychiatric disorders and neuropathic pain syndromes.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481