


Anti-Phospho-STAT3 (Tyr705) Antibody (5W52)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 88 kDa.
Clone:	5W52
Purification:	ProA affinity purified

Applications

Verified Activity: 1. Immunohistochemical analysis of paraffin-embedded human lung tissue using anti-Phospho-STAT3 (Tyr705) antibody. Counter stained with hematoxylin.



Application:	ICC/IF,IHC,IP,WB
Recommended	WB: 1:1000-2000; IHC: 1:50-200; ICC/IF: 1:50-200

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthesized phosphopeptide: human STAT3 around the phosphorylation site of Tyr705
Antigen Species:	human
Uniprot ID:	P40763
Synonyms:	p-STAT3 (Y705);p-STAT3 (Tyr705);STAT3 (p-Tyr705);STAT3 (p-Y705)

Research Background

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of JAK kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α ; and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by Prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481