

Anti-Phospho-PLC γ 1 (Tyr1253) Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Actual: 150 kDa.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Applications

Application:	ELISA,WB
Recommended	WB: 1:500-2000; ELISA: 1:20000

Properties

Stability & Storage:	Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthesized phosphopeptide: human PLC γ 1 around the phosphorylation site of Tyr1253. AA range:1221-1270
Antigen Species:	Human
Uniprot ID:	P19174
Synonyms:	PLC γ 1;PLCgamma1;PLC1;PLC-II;PLC γ 1;PLC γ 1;PLC gamma1;p-PLC γ 1 (Tyr1253);PLC γ 1 (p-Y1253);PLC γ 1;p-PLC γ 1 (Y1253);NCKAP3;IDAA;PLC γ 1 (p-Tyr1253);PLC γ 1;Phospho-PLC γ 1 (Y1253);PLC148;PLC gamma 1;PLC γ 1

Research Background

phospholipase C gamma 1 (PLC γ 1) Homo sapiens The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481