

Anti-USP7 Antibody (8N914)




Product Details



Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Conjugation:	Unconjugated
Molecular Weight:	Theoretical: 128 kDa.
Clone:	8N914
Purification:	ProA affinity purified

Applications

Verified Activity:

1. Western blot analysis of USP7 on 293T cell using anti-USP7 antibody at 1/500 dilution.
2. Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti-USP7 antibody. Counter stained with hematoxylin.
3. Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-USP7 antibody. Counter stained with hematoxylin.
4. Immunohistochemical analysis of paraffin-embedded mouse colon tissue using anti-USP7 antibody. Counter stained with hematoxylin.
5. Flow cytometric analysis of PC-3M cells with USP7 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.





Application: FCM,IHC,WB
Recommended WB: 1:500-1000; IHC: 1:50; FCM: 1:50-100

Properties

Stability & Storage: Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: human USP7 aa 100-200
Antigen Species: Human
Uniprot ID: Q93009
Synonyms: ubiquitin specific peptidase 7 (herpes virus-associated);HAUSP;TEF1

Research Background

HAUSP (herpesvirus-associated ubiquitin-specific protease, USP7) is a ubiquitin-specific protease. HAUSP localizes predominantly to the nucleus, in a TD-dependent manner, where it associates with ND10. ND10 are small nuclear structures implicated in a variety of cellular processes including response to stress and interferons, oncogenesis, and viral infection. HAUSP binds strongly to Vmw110, a herpesvirus regulatory protein which has the ability to disrupt ND10. HAUSP, a novel p53-interacting protein, functions to deubiquitinate p53 in an important pathway for p53 stabilization. HAUSP strongly stabilizes p53 even in the presence of excess Mdm2, and also induces p53-dependent cell growth repression and apoptosis. The HAUSP protein is distributed in the nucleus in a micropunctate pattern with a limited number of larger discrete foci, some of which co-localize with PML in ND10. The gene encoding HAUSP maps to human chromosome band 16p13.2.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481