


GSK717

## Chemical Properties

CAS No. :	1595278-21-9
Formula:	C <sub>28</sub> H <sub>28</sub> N <sub>4</sub> O <sub>2</sub>
Molecular Weight:	452.55
Storage:	Store at low temperature Powder: -20°C for 3 years   In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small>



## Biological Description

Description	<p>GSK717 is a potent, selective NOD2 (nucleotide-binding oligomerization domain 2) inhibitor that inhibits muramyl dipeptide (MDP)-induced NOD2-mediated signaling with an IC<sub>50</sub> of 400 nM for MDP-stimulated IL-8 secretion in HEK293/hNOD2 cells[1]. It blocks the synergy between NOD2 and TLR2 without affecting NOD1, TNFR1, and TLR2-mediated responses. At 5 μM, GSK717 inhibits the release of IL-8, IL-6, TNFα, and IL-1β in primary human monocytes stimulated with MDP[1].</p> <p>[1]. Rickard DJ, et al. Identification of benzimidazole diamides as selective inhibitors of the nucleotide-binding oligomerization domain 2 (NOD2) signaling pathway. PLoS One. 2013;8(8):e69619. Published 2013 Aug 1.</p>
Targets(IC50)	NOD-like Receptor (NLR),NOD,IL Receptor
In vitro	GSK717 (5 μM) inhibits the release of IL-8, IL-6, TNFα and IL-1β in primary human monocytes stimulated with MDP. GSK717 inhibits muramyl dipeptide (MDP)-induced NOD2-mediated signaling, with an IC <sub>50</sub> of 400 nM for MDP-stimulated IL-8 secretion in HEK293/hNOD2 cells[1].

## Solubility Information

Solubility	DMSO: 135 mg/mL (298.31 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	<p>10% DMSO+40% PEG300+5% Tween 80+45% Saline: 5 mg/mL (11.05 mM),Sonication is recommended.</p> <p><i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i></p>

### Preparing Stock Solutions

---

	<b>1mg</b>	<b>5mg</b>	<b>10mg</b>
1 mM	2.2097 mL	11.0485 mL	22.097 mL
5 mM	0.4419 mL	2.2097 mL	4.4194 mL
10 mM	0.221 mL	1.1049 mL	2.2097 mL
50 mM	0.0442 mL	0.221 mL	0.4419 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Rickard DJ, et al. Identification of benzimidazole diamides as selective inhibitors of the nucleotide-binding oligomerization domain 2 (NOD2) signaling pathway. PLoS One. 2013;8(8):e69619. Published 2013 Aug 1.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481