

Rabbit Anti-Human IgG F(ab')₂ Secondary Antibody-Biotin

Product Details

| | |
|---------------|--------------------|
| Ig Type: | IgG |
| Reactivity: | Human |
| Purification: | Protein A purified |

Applications

| | |
|--------------|--|
| Application: | ELISA,ICC/IF,IF,IHC-F,IHC-P,WB |
| Recommended | WB: 1:1000-10000; IHC-P: 1:100-500; IHC-F: 1:100-1000; IF: 1:100-1000; ELISA: 1:1000-10000; ICC/IF: 1:100-1000; FCM: 1:100-1000 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------|--|
| Immunogen: | Human IgG F(ab') ₂ fragment |
| Synonyms: | Rabbit Anti-Human IgG F(ab') ₂ (Biotin) |

Research Background

Human IgG consists of four subclasses (1-4) that can be recognised by antigen differences in their heavy chains. Each subclass has different biological and physiochemical properties. The IgG subclass may be preferentially produced in response to different antigens.

The Fab fragment (fragment antigen binding) is the antigen region of the IgG molecule. It is composed of one constant and one variable domain of each of the heavy and the light chain.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481