


Flusilazole

Chemical Properties

CAS No. :	85509-19-9
Formula:	C ₁₆ H ₁₅ F ₂ N ₃ Si
Molecular Weight:	315.39
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Flusilazole (DPX6573) is an organosilicon fungicide with broad spectrum antifungal effects. Flusilazole can be used to control fungal infections on a variety of vegetables and fruits.
Targets(IC50)	Antifungal

Solubility Information

Solubility	DMSO: 95 mg/mL (301.21 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 3.3 mg/mL (10.46 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.1707 mL	15.8534 mL	31.7068 mL
5 mM	0.6341 mL	3.1707 mL	6.3414 mL
10 mM	0.3171 mL	1.5853 mL	3.1707 mL
50 mM	0.0634 mL	0.3171 mL	0.6341 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

D Fabio Mercado, et al. Reaction Kinetics and Mechanisms of Organosilicon Fungicide Flusilazole With Sulfate and Hydroxyl Radicals. Chemosphere. 2018 Jan;190:327-336.

Ozakca DU, Silah H. Genotoxicity effects of Flusilazole on the somatic cells of Allium cepa. Pestic Biochem Physiol. 2013 Sep;107(1):38-43.

Dimopoulou M, et al. Flusilazole induces spatio-temporal expression patterns of retinoic acid-, differentiation- and sterol biosynthesis-related genes in the rat Whole Embryo Culture. Reprod Toxicol. 2016 Sep;64:77-85.

Yu S, et al. Residue and dissipation dynamics of flusilazole in apple and soil. Bull Environ Contam Toxicol. 2011 Mar;86(3):319-22.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481