

CD98 Protein, Mouse, Recombinant (His) V2

General Information

| | |
|-----------------------|---|
| Synonyms: | NACAE;solute carrier family 3 (amino acid transporter heavy chain), member 2;AI314110;Ly10;Cd98;4F2;Ly-m10;Mgp-2hc;4F2HC;Mdu1;Ly-10 |
| Protein Construction: | Ala100-Ala526 |
| Species: | Mouse |
| Expression Host: | HEK293 Cells |
| Accession: | P10852-1 |
| Molecular Weight: | 48.67 kDa (Predicted); 55-75 kDa (Due to glycosylation) |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. |
| Purity: | > 95% as determined by Bis-Tris PAGE; > 95% as determined by HPLC |
| Endotoxin: | < 1.0 EU/ μ g of the protein as determined by the LAL method. |
| Formulation: | Lyophilized from 0.22 μ m filtered solution in PBS, 200 mM L-arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μ g/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C . For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The type II transmembrane protein CD98, best known as the heavy chain of the heterodimeric amino acid transporters (HAT), is required for the surface expression and basolateral localization of this transporter complex in polarized epithelial cells. CD98 also interacts with beta1 integrins resulting in an increase in their affinity for ligand. In this study we explored the role of the transmembrane and cytoplasmic domains of CD98 on integrin-dependent cell adhesion and migration in polarized renal epithelial cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481