

TFF1 Protein, Human, Recombinant (6xHis)

General Information

Synonyms:	HPS2;D21S21;pNR-2;trefoil factor 1;HP1.A;BCEI;pS2
Protein Construction:	Glu25-Phe84
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P04155
Molecular Weight:	7.5 kDa (Predicted); 14 kDa (Reducing conditions)

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it.
Purity:	> 95% as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in sterile deionized water. The product concentration should not be less than 100 μg/mL. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Trefoil Factor 1 (TFF1) belongs to the three structurally related secreted proteins that contain trefoil domains. TFF1 is an approximately 7 kDa peptide that plays an important role in epithelial regeneration and wound healing. It is highly expressed in goblet cells of the gastric and intestinal mucosa and by conjunctival goblet cells. By conserving intrachain disulfide bonds, human TFF1 formed a three-leaved conformation held together. It is a copper-binding protein that can form disulfide-linked homodimers, associate into disulfide-linked complexes with Gastrokine 2, and form non-covalent complexes with the mucin MUC5AC. TFF1 is down-regulated during the progression from gastritis to gastric dysplasia to gastric cancer, although it is up-regulated in breast and prostate

cancers.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481