

Anti-CD15/SSEA1 Antibody (3F592)

Product Details

Ig Type:	IgM/Kappa
Reactivity:	Human,Rat
Clone:	3F592
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Rat Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Cervical Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Endometrial Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Human Lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
10. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal

A DRUG SCREENING EXPERT

Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Placenta; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD15/SSEA1 Monoclonal Antibody, Unconjugated (TMAB-03856) at 1:100 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.

Application: IHC-P,IHC-Fr,IF,FCM,ICC/IF

Recommended IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1:100-200; ICC/IF: 1:100-500

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated synthetic peptide: human CD15

Antigen Species: Human

Gene ID: 2526

Uniprot ID: P22083

Research Background

The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metastasis, inflammation, and bacterial adhesion. They are secondarily absorbed to red blood cells giving rise to their Lewis phenotype. This gene is a member of the fucosyltransferase family, which catalyzes the addition of fucose to precursor polysaccharides in the last step of Lewis antigen biosynthesis. It encodes an enzyme with alpha(1,3)-fucosyltransferase and alpha(1,4)-fucosyltransferase activities. Mutations in this gene are responsible for the majority of Lewis antigen-negative phenotypes. Multiple alternatively spliced variants, encoding the same protein, have been found for this gene. [provided by RefSeq].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481