

Anti-CD162 Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human
Molecular Weight:	Theoretical: 41 kDa. Actual: 41 kDa.
Purification:	Protein A purified

Applications

1. Sample:

Raji (Human) Cell Lysate at 30 µg

U937 (Human) Cell Lysate at 30 µg

Primary: Anti-CD162 (TMAB-03865) at 1/500 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 41 kD

Observed band size: 41 kD

2. Sample:

Jurkat (Human) Cell Lysate at 30 µg

MOLT-4 (Human) Cell Lysate at 30 µg

Primary: Anti-CD162 (TMAB-03865) at 1/500 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 41 kD

Observed band size: 41 kD

3. Blank control: U937 (blue).

Verified Activity: Primary Antibody: Rabbit Anti-CD162 antibody (TMAB-03865), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG (orange), used under the same conditions.

Secondary Antibody: Goat anti-rabbit IgG-PE (white blue), Dilution: 1: 200 in 1 X PBS containing 0.5% BSA.

Protocol

The cells were fixed with 2% paraformaldehyde (10 min). Primary antibody (TMAB-03865, 1 µg /1x10⁶ cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA+10% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Anti-rabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.

4. MCF7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Antibody incubation with (CD162) polyclonal Antibody, Unconjugated (TMAB-03865) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue) was used to stain the cell nuclei.

A DRUG SCREENING EXPERT

Application: WB,ICC/IF,FCM

Recommended WB: 1:500-2000; ICC/IF: 1:50-200; FCM: 1µg/Test

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated synthetic peptide: human PSGL-1

Antigen Species: Human

Gene ID: 6404

Uniprot ID: Q14242

Research Background

This gene encodes a glycoprotein that functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Apr 2011].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481