

Anti-CD27 Polyclonal Antibody 2

Product Details

Ig Type:	IgG
Reactivity:	Mouse,Rat (predicted:Human,Dog,Pig,Cow)
Molecular Weight:	Theoretical: 27 kDa. Actual: 60 kDa.
Purification:	Protein A purified

Applications

Sample:

Lane 1: Lymph node (Mouse) Lysate at 30 µg

Lane 2: Thymus (Rat) Lysate at 40 µg

Lane 3: Spleen (Rat) Lysate at 40 µg

Lane 4: Lymph node (Rat) Lysate at 30 µg

Verified Activity:

Primary:

Anti-CD27 (TMAB-03927) at 1/1000 dilution

Anti-beta-Actin at 1/2000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 50-55 kD

Observed band size: 60 kD

Application:

WB

Recommended

WB: 1:500-2000

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated synthetic peptide: human CD27
Antigen Species:	Human
Gene ID:	939
Uniprot ID:	P26842

Research Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor. [provided by RefSeq, Jul 2008]

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481