

## Anti-Collagen III Antibody (5Y376)

### Product Details

Ig Type:	IgG
Reactivity:	Human
Molecular Weight:	Theoretical: 117 kDa. Actual: 150-200 kDa.
Clone:	5Y376
Purification:	Protein G purified

### Applications

#### Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Human Esophagus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Prostate Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Abdominal skin; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen III Monoclonal Antibody, Unconjugated (TMAB-04523) at 1:1000 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.
9. 25 µg total protein per lane of various lysates (see on figure) probed with Collagen III monoclonal antibody, unconjugated (TMAB-04523) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
10. 4% Paraformaldehyde-fixed RD (H) cell; Triton X-100 at r. T. for 20 min; Antibody incubation

## A DRUG SCREENING EXPERT

---

with (Collagen III) monoclonal Antibody, unconjugated (TMAB-04523) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Mouse IgG antibody (green) at 37°C for 90 min, DAPI (blue) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

Application: WB,IHC-P,IHC-Fr,ICC/IF,IF

Recommended WB: 1:500-2000; IHC-P: 1:500-1000; IHC-Fr: 1:500-1000; ICC/IF: 1:100-200; IF: 1:500-1000

---

### Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

---

### Antigen Details

Immunogen: Recombinant Protein: human Collagen III protein

Antigen Species: Human

Gene ID: 1281

Uniprot ID: P02461

---

### Research Background

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function.

This gene encodes the pro-alpha1 chains of type III collagen, a fibrillar collagen that is found in extensible connective tissues such as skin, lung, uterus, intestine and the vascular system, frequently in association with type I collagen. Mutations in this gene are associated with Ehlers-Danlos syndrome types IV, and with aortic and arterial aneurysms. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481

---