

Anti-Fbxw7 Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat (predicted:Dog,Pig,Cow,Horse)
Molecular Weight:	Theoretical: 78 kDa.
Purification:	Protein A purified

Applications

1. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-Fbxw7/CDC4 Polyclonal Antibody, Unconjugated (TMAB-05938) 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining

2. Blank control: Mouse spleen.

Primary Antibody (green line): Rabbit Anti-Fbxw7 antibody (TMAB-05938)

Dilution: 2 µg /10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG.

Secondary Antibody: Goat anti-rabbit IgG-FITC

Dilution: 1 µg /test.

Protocol

The cells were fixed with 4% PFA (10 min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Verified Activity:

3. Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Fbxw7) Polyclonal Antibody, Unconjugated (TMAB-05938) at 1:400 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

4. Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Fbxw7) Polyclonal Antibody, Unconjugated (TMAB-05938) at 1:400 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.

5. Blank control (black line): MCF-7.

Primary Antibody (green line): Rabbit Anti-Fbxw7 antibody (TMAB-05938)

Dilution: 1 µg/Test;

Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488

Dilution: 0.5 µg/Test.

Isotype control (orange line): Normal Rabbit IgG

Protocol

The cells were fixed with 4% PFA (10 min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5% BSA to block

non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

6. Blank control (black line): MCF-7.

Primary Antibody (green line): Rabbit Anti-Fbxw7 antibody (TMAB-05938)

Dilution: 1 µg/Test;

Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488

Dilution: 0.5 µg/Test.

Isotype control (orange line): Normal Rabbit IgG

Protocol

The cells were fixed with 4% PFA (10 min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Application: IHC-P,IHC-Fr,IF,FCM

Recommended IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1µg/Test

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated synthetic peptide: human Fbxw7/CDC4

Antigen Species: Human

Gene ID: 55294

Uniprot ID: Q969H0

Research Background

Fbw7 is a member of the F box protein family which are characterized by an approximately 40 amino acid motif, the F box. The F box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F box), which function in phosphorylation-dependent ubiquitination. The F box proteins are divided into 3 classes: Fbws containing WD40 domains, Fbls containing leucine rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. Fbw7 belongs to the Fbws class; in addition to an F box, this protein contains 7 tandem WD40 repeats. It binds directly to cyclin E and probably targets cyclin E for ubiquitin mediated degradation. Mutations of this gene are detected in ovarian and breast cancer cell lines. Alternative splicing of this gene generates 2 transcript variants diverging at the 5' termini.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481