

Anti-Geminin Antibody (1B890)

Product Details

Ig Type:	IgG
Reactivity:	Human
Molecular Weight:	Theoretical: 24 kDa. Actual: 27 kDa.
Clone:	1B890
Purification:	Protein A purified

Applications

Verified Activity:	1. Paraformaldehyde-fixed, paraffin embedded Human Ovarian Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation Geminin Monoclonal Antibody, Unconjugated (TMAB-06435) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
	2. Paraformaldehyde-fixed, paraffin embedded Human Esophagus Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation Geminin Monoclonal Antibody, Unconjugated (TMAB-06435) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
	3. 4% Paraformaldehyde-fixed HepG2 (H) cell; Triton X-100 at r. T. for 20 min; Antibody incubation with (Geminin) monoclonal Antibody, unconjugated (TMAB-06435) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green) at 37°C for 90 min, DAPI (blue) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.
	4. 25 µg total protein per lane of various lysates (see on figure) probed with Geminin monoclonal antibody, unconjugated (TMAB-06435) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
Application:	WB,IHC-P,IHC-Fr,ICC/IF,IF
Recommended	WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:50-200; IF: 1:100-500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthesized peptide: human Geminin
Antigen Species:	Human
Gene ID:	51053
Uniprot ID:	O75496

Research Background

Inhibits DNA replication by preventing the incorporation of MCM complex into prereplication complex (pre-RC). It is degraded during the mitotic phase of the cell cycle. Its destruction at the metaphase-anaphase transition permits

replication in the succeeding cell cycle.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481