

Anti-GLEPP1 Polyclonal Antibody

Product Details

| | |
|-------------------|---------------------------------------|
| Ig Type: | IgG |
| Reactivity: | Rat (predicted:Human,Mouse,Dog,Horse) |
| Molecular Weight: | Theoretical: 135 kDa. |
| Purification: | Protein A purified |

Applications

| | |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Verified Activity: | Tissue/cell: rat lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-GGA2 Polyclonal Antibody, Unconjugated 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining |
| Application: | IHC-P,IHC-Fr,IF |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

| | |
|----------------------|---------------------------------------------------------------------------------------------------------|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|------------------------------------------------|
| Immunogen: | KLH conjugated synthetic peptide: human GLEPP1 |
| Antigen Species: | Human |
| Gene ID: | 5800 |
| Uniprot ID: | Q16827 |

Research Background

Protein phosphatases play critical roles in the regulation of signal transduction pathways. The family can be separated into three general categories, which are distinguished on the basis of substrate specificity. The first and largest category, termed protein tyrosine phosphatases or PTPs, includes transmembrane proteins, membrane associated proteins and proteins that localize to nuclei. The second category of protein phosphatases dephosphorylate proteins on phosphoserine and phosphothreonine residues, whereas the third category of protein phosphatases exhibit dual specificities and can dephosphorylate proteins on phosphotyrosine and phosphoserine/phosphothreonine residues. Glomerular epithelial protein 1 (GLEPP1), also designated protein tyrosine phosphatase U2 (PTPase U2) or receptor-type tyrosine-protein phosphatase O (PTPRO), belongs to the protein-tyrosine phosphatase family. GLEPP1 is a Type I membrane protein containing 8 fibronectin type-III domains and 1 tyrosine-protein phosphatase domain. GLEPP1 is expressed primarily in the glomeruli of the kidney, but is also detected in placenta, lung and brain.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481