

Anti-Glutaredoxin 2 Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human,Rat (predicted:Mouse,Dog,Cow,Rabbit,Sheep)
Molecular Weight:	Theoretical: 16 kDa.
Purification:	Protein A purified

Applications

Verified Activity:	<p>1. Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-GLRX2 Polyclonal Antibody, Unconjugated (TMAB-06562) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining</p> <p>2. Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (GLRX2) Polyclonal Antibody, Unconjugated (TMAB-06562) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.</p> <p>3. Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (GLRX2) Polyclonal Antibody, Unconjugated (TMAB-06562) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.</p> <p>4. Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (GLRX2) Polyclonal Antibody, Unconjugated (TMAB-06562) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.</p> <p>5. Blank control (blue line): RSC96 (fixed with pre-warmed 4% paraformaldehyde for 30 min at 37°C and then permeabilized with 90% ice-cold methanol for 30 min on ice) Primary Antibody (green line): Rabbit Anti-Glutaredoxin 2 antibody (TMAB-06562), Dilution: 0.2 µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE, Dilution: 1 µg /test.</p>
Application:	IHC-P,IHC-Fr,IF,FCM
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1µg/Test

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: KLH conjugated synthetic peptide: human Glutaredoxin 2

Antigen Species: Human

Gene ID: 51022

Uniprot ID: Q9NS18

Research Background

Glutathione-dependent oxidoreductase that facilitates the maintenance of mitochondrial redox homeostasis upon induction of apoptosis by oxidative stress. Involved in response to hydrogen peroxide and regulation of apoptosis caused by oxidative stress. Acts as a very efficient catalyst of monothiol reactions because of its high affinity for protein glutathione-mixed disulfides. Can receive electrons not only from glutathione (GSH), but also from thioredoxin reductase supporting both monothiol and dithiol reactions. Efficiently catalyzes both glutathionylation and deglutathionylation of mitochondrial complex I, which in turn regulates the superoxide production by the complex. Overexpression decreases the susceptibility to apoptosis and prevents loss of cardiolipin and cytochrome c release.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481