

Anti-GRB 14 Polyclonal Antibody

Product Details

| | |
|-------------------|--|
| Ig Type: | IgG |
| Reactivity: | Human (predicted:Mouse,Rat,Chicken,Dog,Cow,Horse,Rabbit,Sheep) |
| Molecular Weight: | Theoretical: 61 kDa. |
| Purification: | Protein A purified |

Applications

| | |
|--------------------|---|
| Verified Activity: | Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-GRB14 Polyclonal Antibody, Unconjugated (TMAB-06761) 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining |
| Application: | IHC-P,IHC-Fr,IF |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | KLH conjugated synthetic peptide: human GRB 14 |
| Antigen Species: | Human |
| Gene ID: | 2888 |
| Uniprot ID: | Q14449 |

Research Background

Many growth factors function by binding receptors with intrinsic tyrosine kinase activity. Signaling by such receptors involves a series of intermediates characterized by SH2 domains that bind tyrosine phosphorylated receptors by a direct interaction between the SH2 domain and specific phospho-tyrosine-containing receptor sequences. GRB7, a SH2 domain protein, has a single SH2 domain at its C-terminal, a central region with similarity to Ras GAP and a proline-rich N-terminus. A related SH2 domain-containing protein, GRB10, exhibits a high degree of homology with GRB7. GRB10 undergoes serine but not tyrosine phosphorylation in response to EGF treatment, but appears to bind to the EGF receptor poorly. An additional member of the GRB7 family, designated GRB14, contains a Pleckstrin homology domain in its central region and a carboxy-terminal SH2 domain. GRB14 mRNA is expressed at high levels in a broad range of tissues including liver, kidney, pancreas, testis, ovary, heart and skeletal muscle. Expression of the GRB14 protein in breast carcinomas is strongly correlated with estrogen receptor positivity.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481