

## Anti-HCN2 Polyclonal Antibody 2

## Product Details

Ig Type:	IgG
Reactivity:	Human (predicted:Mouse,Rat,Dog,Cow,Rabbit,Sheep)
Molecular Weight:	Theoretical: 97 kDa. Actual: 110 kDa.
Purification:	Protein A purified

## Applications

Sample:	Lane 1: Human Jurkat cell lysates Lane 2: Human SH-SY5Y cell lysates
Verified Activity:	Primary: Anti-HCN2 (TMAB-06916) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 97 kDa Observed band size: 110 kDa
Application:	WB
Recommended	WB: 1:500-2000

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	KLH conjugated synthetic peptide: human HCN2
Antigen Species:	Human
Gene ID:	610
Uniprot ID:	Q9UL51

## Research Background

Hyperpolarization-activated, cyclic nucleotide-binding channels (HCN) are voltage-gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family consists of four members (HCN1-4), each with a core transmembrane segment domain and a carboxy-terminal 120 amino-acid cyclic nucleotide-binding domain motif (1). HCN channels are expressed in the brain, heart, thalamus and testis (1). The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart (1). The genes encoding human HCN1 and HCN2 map to chromosomes 5 and 19p13.3, respectively (2,3). The genes encoding HCN3 and HCN4 map to chromosomes 1q21.3 and 15q24-q25, respectively (4,5).

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481