

Anti-Acetyl-Histone H2B (Lys5) Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Human (predicted:Mouse,Rat)
Molecular Weight:	Theoretical: 14 kDa.
Purification:	Protein A purified

Applications

	Blocking buffer: 5% NFDM/TBST	
	Primary ab dilution: 1: 2000	
	Primary ab incubation condition: 2 hours at	room temperature
	Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)	
Verified Activity:	Lysate: (-) HeLa, (+) HeLa+Sodium butyrate	(30 mM, 4 h)
	Protein loading quantity: 20 µg	
	Exposure time: 60 s	
	Predicted MW: 14 kDa	
	Observed MW: 14 kDa	
Application:	WB,IHC-P,IHC-Fr,IF	
Recommended	WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500	

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated synthetic peptide: human Histone H2B (Acetyl K5)
Antigen Species:	Human
Gene ID:	3018
Uniprot ID:	P33778

Research Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481