

## Anti-Histone H2B/HIST1H2BD Polyclonal Antibody

### Product Details

Ig Type:	IgG
Reactivity:	Human,Rat (predicted:Mouse,Chicken,Dog,Pig,Cow,Horse,Rabbit)
Molecular Weight:	Theoretical: 14 kDa.
Purification:	Protein A purified

### Applications

Verified Activity:	<p>1. Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Histone H2B/HIST1 h2BD) Polyclonal Antibody, Unconjugated (TMAB-07092) at 1: 200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.</p> <p>2. Blank control (black line): Hela. Primary Antibody (green line): Rabbit Anti-HIST1 h2BD antibody (TMAB-07092) Dilution: 1 µg/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5 µg/Test. Isotype control (orange line): Normal Rabbit IgG</p>
	<p>Protocol The cells were fixed with 4% PFA (10 min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.</p>
Application:	IHC-P,IHC-Fr,IF,FCM
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1µg/Test

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen: KLH conjugated synthetic peptide: human Histone H2B  
Antigen Species: Human  
Gene ID: 3018  
Uniprot ID: P33778

---

### Research Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481