

Anti-HPGD Antibody (80345)

Product Details

Ig Type:	IgG
Reactivity:	Human
Molecular Weight:	Theoretical: 29 kDa. Actual: 24 kDa.
Clone:	80345
Purification:	Protein A purified

Applications

Verified Activity:	<p>1. 25 ug total protein per lane of various lysates (see on figure) probed with HPGD monoclonal antibody, unconjugated (TMAB-07274) at 1: 1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.</p> <p>2. The A549 (H) cells were fixed with 4% PFA (10 min at r. T.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r. T.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green): Rabbit Anti-HPGD antibody (TMAB-07274, 1: 100); Isotype Control (orange): Rabbit IgG. Blank control (black): PBS. Acquisition of 20,000 events was performed.</p>
Application:	FCM,WB
Recommended	FCM=1:50-100; WB=1:500-2000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	A synthesized peptide: human 15 PGDH
Antigen Species:	Human
Gene ID:	3248
Uniprot ID:	P15428

Research Background

Catalyzes the NAD-dependent dehydrogenation (oxidation) of a broad array of hydroxylated polyunsaturated fatty acids (mainly eicosanoids and docosanoids, including prostaglandins, lipoxins and resolvins), yielding their corresponding keto (oxo) metabolites.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481