

Anti-human CD18 Antibody (9C890)

Product Details

Ig Type:	Mouse IgG1, k
Reactivity:	Human
Molecular Weight:	Theoretical: 82 kDa. Actual: 90-95 kDa.
Clone:	9C890
Purification:	Protein G purified

Applications

Verified Activity:	<p>1. The human blood granulocytes (H) cells were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r. T.). Primary Antibody (green): Mouse Anti-CD18 antibody (TMAB-07389, 1:100); Isotype Control (orange): Mouse IgG. Blank control (black): PBS. Acquisition of 20,000 events was performed.</p> <p>2. Paraformaldehyde-fixed, paraffin embedded Human Lung; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with human CD18 Monoclonal Antibody, Unconjugated (TMAB-07389) at 1: 200 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.</p> <p>3. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with human CD18 Monoclonal Antibody, Unconjugated (TMAB-07389) at 1: 200 overnight at 4°C, followed by conjugation to the AffiniPure Goat Anti-Mouse IgG H&L-HRP and DAB staining.</p>
Application:	IHC-P,IHC-Fr,IF,FCM
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500; FCM: 1:50-100

Properties

Stability & Storage:	Store at 2°C-8°C for 12 months, do not freeze. Keep away from direct sunlight.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated synthetic peptide: human CD18
Antigen Species:	Human
Gene ID:	3689
Uniprot ID:	P05107

Research Background

The product of this gene belongs to the integrin beta chain family of proteins. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. This gene encodes the integrin beta chain beta 2. A given chain may combine with multiple partners resulting in different integrins. For example, beta 2 combines with the alpha L chain to form the integrin LFA-1, and combines with the alpha M chain to form the integrin Mac-1. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling. Defects in this gene are the cause of leukocyte adhesion deficiency type I (LAD1). Two transcript variants encoding the same protein have been

identified for this gene. [provided by RefSeq, Jul 2008].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481