

## Anti-JNK1 Antibody (8M452)

## Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 48 kDa. Actual: 43,50 kDa.
Clone:	8M452
Purification:	Protein A purified

## Applications

Verified Activity:	<p>1. 25 µg total protein per lane of various lysates (see on figure) probed with JNK1 monoclonal antibody, unconjugated (TMAB-07879) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.</p> <p>2. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JNK1 Monoclonal Antibody, Unconjugated (TMAB-07879) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&amp;L Secondary Antibody-HRP and DAB staining.</p> <p>3. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JNK1 Monoclonal Antibody, Unconjugated (TMAB-07879) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&amp;L Secondary Antibody-HRP and DAB staining.</p> <p>4. Paraformaldehyde-fixed, paraffin embedded Human Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JNK1 Monoclonal Antibody, Unconjugated (TMAB-07879) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&amp;L Secondary Antibody-HRP and DAB staining.</p>
Application:	WB,IHC-P,IHC-Fr,IF
Recommended	WB: 1:500-2000; IHC-P: 1:50-200; IHC-Fr: 1:50-200; IF: 1:50-200

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	A synthesized peptide: human JNK1
Antigen Species:	Human
Gene ID:	5599
Uniprot ID:	P45983

## Research Background

phosphorylated at the Thr-Pro-Tyr phosphorylation motif instead of the characteristic MAP kinase Thr-Glu-Tyr motif. JNK2 (p54a, SAPK1a), along with JNK1 and JNK3, is thought to play an important role in nuclear signal transduction through its environmental stress activation and subsequent phosphorylation of the nuclear transcription factor p53.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481