

Anti-MAPT/Tau Antibody (1F539)

Product Details

Ig Type:	IgG3
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 46/79 kDa. Actual: 50-70 kDa.
Clone:	1F539
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Human Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Human Breast; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Human Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Human Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Human Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
5. 25 ug total protein per lane of various lysates (see on figure) probed with Tau-4 monoclonal antibody, unconjugated (TMAB-08598) at 1: 1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
6. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Tau-4 Monoclonal Antibody, Unconjugated (ascites of TMAB-08598) at 1: 1500 overnight at 4°C, followed by conjugation to the HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Tau-4 Monoclonal Antibody, Unconjugated (ascites of TMAB-08598) at 1: 1500 overnight at 4°C, followed by conjugation to the HRP and DAB staining.
8. (Negative control) Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Tau-4 Monoclonal Antibody, Unconjugated (ascites of TMAB-08598) at 1: 1500 overnight at 4°C, followed by conjugation to the HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MAPT/Tau Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (Red, BF594), DAPI (blue) was used to stain the cell nuclei.
10. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in

A DRUG SCREENING EXPERT

sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MAPT/Tau Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (Red, BF594), DAPI (blue) was used to stain the cell nuclei.

11. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MAPT/Tau Monoclonal Antibody, Unconjugated (TMAB-08598) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (Red, BF594), DAPI (blue) was used to stain the cell nuclei.

Application: WB,IHC-P,IHC-Fr,IF

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: Recombinant Protein: Human Tau-4 Protein

Antigen Species: Human

Gene ID: 4137

Uniprot ID: P10636-8

Research Background

Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481
