

## Anti-MAPT/Tau Antibody (4Q63)

### Product Details

Ig Type:	IgG1
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 46/79 kDa. Actual: 50-70 kDa.
Clone:	4Q63
Purification:	Protein A purified

### Applications

1. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Breast; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Left Parietal Lobe; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C, followed by conjugation to the SP Kit (Mouse) and DAB staining.
7. 25 µg total protein per lane of various lysates (see on figure) probed with Tau-4 monoclonal antibody, unconjugated (TMAB-08600) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
8. Paraformaldehyde-fixed, paraffin embedded Human Breast; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Tau-4 Monoclonal Antibody, Unconjugated (TMAB-08600) at 1: 200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (green), DAPI (blue) was used to stain the cell nuclei.

Verified Activity:

## A DRUG SCREENING EXPERT

---

Application: IHC-P,IHC-Fr,IF,WB

Recommended IHC-P: 1:200-500; IHC-Fr: 1:200-500; IF: 1:200-500; WB: 1:500-2000

---

### Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

---

### Antigen Details

Immunogen: Recombinant Protein: human Tau-4 protein

Antigen Species: Human

Gene ID: 4137

Uniprot ID: P10636

---

### Research Background

Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481

---