

Anti-MMP1 Antibody (2G245)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse
Molecular Weight:	Theoretical: 54 kDa. Actual: 54 kDa.
Clone:	2G245
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Mouse Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Cervical Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Placenta; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Esophagus Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with MMP1 Monoclonal Antibody, Unconjugated (TMAB-08858) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

A DRUG SCREENING EXPERT

Application: IHC-P,IHC-Fr,ICC/IF,IF,FCM

Recommended IHC-P: 1:50-200; IHC-Fr: 1:50-200; ICC/IF: 1:50-200; IF: 1:50-200; FCM: 1:50-100

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human MMP1

Antigen Species: Human

Gene ID: 4312

Uniprot ID: P03956

Research Background

Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X. In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481