

Anti-MOGT1 Polyclonal Antibody

Product Details

| | |
|-------------------|--|
| Ig Type: | IgG |
| Reactivity: | Mouse (predicted:Human,Rat,Pig,Rabbit) |
| Molecular Weight: | Theoretical: 39 kDa. |
| Purification: | Protein A purified |

Applications

| | |
|--------------------|---|
| Verified Activity: | Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (MOGT1) Polyclonal Antibody, Unconjugated (TMAB-08904) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining. |
| Application: | IHC-P,IHC-Fr,IF |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|---|
| Immunogen: | KLH conjugated synthetic peptide: human MOGT1 |
| Antigen Species: | Human |
| Gene ID: | 116255 |
| Uniprot ID: | Q96PD6 |

Research Background

Acyl-CoA:monoacylglycerol acyltransferase (MOGAT; EC 2.3.1.22) catalyzes the synthesis of diacylglycerols, the precursor of physiologically important lipids such as triacylglycerol and phospholipids (Yen et al., 2002 [PubMed 12077311]).[supplied by OMIM, Mar 2008]

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481