

Anti-NAP2 Polyclonal Antibody 2

Product Details

| | |
|-------------------|---------------------------------|
| Ig Type: | IgG |
| Reactivity: | Human (predicted:Mouse,Rat,Pig) |
| Molecular Weight: | Theoretical: 14 kDa. |
| Purification: | Protein A purified |

Applications

| | |
|--------------------|--|
| Verified Activity: | Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-NAP2 Polyclonal Antibody, Unconjugated (TMAB-09234) 1: 200, overnight at 4° C, followed by conjugation to the secondary antibody and DAB staining |
| Application: | IHC-P,IHC-Fr,IF |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|---|
| Immunogen: | KLH conjugated synthetic peptide: mouse CXCL7 |
| Antigen Species: | Mouse |
| Gene ID: | 57349 |

Research Background

The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. [provided by RefSeq, May 2010].

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481