

## Anti-Netrin G1 ligand Polyclonal Antibody

### Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat (predicted:Dog,Cow,Horse,Rabbit)
Molecular Weight:	Theoretical: 70 kDa. Actual: 76 kDa.
Purification:	Protein A purified

### Applications

Verified Activity:	1. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-NGL1 Polyclonal Antibody, Unconjugated (TMAB-09368) 1: 200, overnight at 4° C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated used at 1: 200 dilution for 40 minutes at 37°C. DAPI (5 µg/ml, blue) was used to stain the cell nuclei
	2. Tissue/cell: mouse brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-NGL1 Polyclonal Antibody, Unconjugated (TMAB-09368) 1: 200, overnight at 4° C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated used at 1: 200 dilution for 40 minutes at 37°C. DAPI (5 µg/ml, blue) was used to stain the cell nuclei
	3. Sample: Hela Cell (Human) Lysate at 40 µg Primary: Anti-Netrin G1 ligand (TMAB-09368) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 76 kD
	4. Sample: BV2 Cell (Mouse) Lysate at 40 µg Primary: Anti-Netrin G1 ligand (TMAB-09368) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 70 kD Observed band size: 76 kD
Application:	WB,IHC-P,IHC-Fr,IF
Recommended	WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen: KLH conjugated synthetic peptide: human Netrin G1 ligand  
Antigen Species: Human  
Gene ID: 57689  
Uniprot ID: Q9HCJ2

---

### Research Background

NGL-1 is a single pass type I membrane protein that acts as a cell adhesion molecule. It contains nine leucine-rich repeats (LRR) and one Ig-like C2-type domain. NGL-1 is predominantly expressed in the striatum and the cerebral cortex of both the embryonic and adult brain. NGL-1 specifically interacts with Netrin G1 (a molecule involved in axon guidance in the developing central nervous system) via its LRR region. NGL-1 plays a role in the regulation of neurite outgrowth of developing thalamic neurons. Soluble NGL-1 inhibits thalamic axon outgrowth while NGL-1 that is bound to the surface of developing thalamocortical axons stimulates growth. NGL-1 also interacts with Whirlin possibly stabilizing interstereociliar links.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481