

Anti-NMT55 Antibody (2E849)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 54 kDa. Actual: 63 kDa.
Clone:	2E849
Purification:	Protein A purified

Applications

Verified Activity:

1. The A673 (H) cells were fixed with 4% PFA (10 min at r. T.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r. T.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green): Rabbit Anti-NMT55 antibody (TMAB-09541, 1: 100); Isotype Control (orange): Rabbit IgG. Blank control (black): PBS. Acquisition of 20,000 events was performed.
2. 25 ug total protein per lane of various lysates (see on figure) probed with NMT55 monoclonal antibody, unconjugated (TMAB-09541) at 1: 1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
3. Paraformaldehyde-fixed, paraffin embedded Mouse Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Rat Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Mouse Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Rat Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to

the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

10. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Mouse Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

12. Paraformaldehyde-fixed, paraffin embedded Rat Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

13. Paraformaldehyde-fixed, paraffin embedded Human Prostate; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

14. Paraformaldehyde-fixed, paraffin embedded Mouse Prostate; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with NMT55 Monoclonal Antibody, Unconjugated (TMAB-09541) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

Application: FCM,ICC/IF,IF,IHC-Fr,IHC-P,WB

Recommended FCM=1:50-100; ICC/IF=1:50-200; IF=1:100-500; IHC-Fr=1:100-500; IHC-P=1:100-500; WB=1:500-2000

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human NONO

Antigen Species: Human

Gene ID: 4841

Uniprot ID: Q15233

Research Background

DNA- and RNA binding protein, involved in several nuclear processes. Binds the conventional octamer sequence in double-stranded DNA. Also binds single-stranded DNA and RNA at a site independent of the duplex site. Involved in pre-mRNA splicing, probably as a heterodimer with SFPQ.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481