

Anti-PDHB Antibody (7G981)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 39 kDa. Actual: 37 kDa.
Clone:	7G981
Purification:	Protein A purified

Applications

Verified Activity:

1. 25 µg total protein per lane of various lysates (see on figure) probed with PDHB monoclonal antibody, unconjugated (TMAB-10127) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
2. Paraformaldehyde-fixed, paraffin embedded Mouse Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Rat Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Rat Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Rat Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
10. Paraformaldehyde-fixed, paraffin embedded Human Pancreas; Antigen retrieval by boiling

in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

12. Paraformaldehyde-fixed, paraffin embedded Human Thyroid Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

13. Paraformaldehyde-fixed, paraffin embedded Human Ovarian Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

14. Paraformaldehyde-fixed, paraffin embedded Human Gastric Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PDHB Monoclonal Antibody, Unconjugated (TMAB-10127) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

Application: WB,IHC-P,IHC-Fr,ICC/IF,IF,IP,FCM

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:50-200; IF: 1:100-500; IP: 1:20-50; FCM: 1:50-100

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human PDHB

Antigen Species: Human

Gene ID: 5162

Uniprot ID: P11177

Research Background

The pyruvate dehydrogenase complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂, and thereby links the glycolytic pathway to the tricarboxylic cycle.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481