

## Anti-PDPK1 Antibody (1V774)

### Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse
Molecular Weight:	Theoretical: 61 kDa.
Clone:	1V774
Purification:	Protein A purified

### Applications

Verified Activity:	1. Cell line: NIH3T3
	Fixative: 4% Paraformaldehyde
	Permeabilization: 0.1% TritonX-100
	Primary ab dilution: 1:50
	Primary incubation condition: 4°C overnight
	Secondary ab: Goat Anti-Rabbit IgG
	Nuclear counter stain: DAPI (Blue)
	Comment: Color green is the positive signal for TMAB-10148
	2. Tissue: Human liver cancer
	Section type: Formalin fixed & Paraffin-embedded section
	Retrieval method: High temperature and high pressure
	Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100
	Primary ab incubation condition: 1 hour at room temperature
	Counter stain: Hematoxylin
	Comment: Color brown is the positive signal for TMAB-10148
3. Tissue: Human liver	
Section type: Formalin fixed & Paraffin-embedded section	
Retrieval method: High temperature and high pressure	
Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100	
Primary ab incubation condition: 1 hour at room temperature	
Counter stain: Hematoxylin	
Comment: Color brown is the positive signal for TMAB-10148	
Application:	IHC-P,IHC-Fr,ICC/IF,IF
Recommended	IHC-P: 1:50-200; IHC-Fr: 1:50-200; ICC/IF: 1:50-200; IF: 1:50-200

### Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

### Antigen Details

Immunogen: KLH conjugated synthetic peptide: human PDPK1  
Antigen Species: Human  
Gene ID: 5170  
Uniprot ID: O15530

---

### Research Background

PDK1 (3 Phosphoinositide Dependent Protein Kinase 1) phosphorylates AGC kinases. PDK1 activates conventional PKC and PKC zeta through phosphorylation of critical threonine residues in the activation loop. PDK1 also phosphorylates Protein Kinase B (PKB) at threonine 308 in the presence of phosphatidylinositol-3,4,5-trisphosphate. Active Akt inactivates Glycogen Synthase Kinase 3 (GSK3), eventually leading to the dephosphorylation and activation of glycogen synthase and the stimulation of glycogen synthesis. Because of the role that PDK plays in insulin-induced glycogen synthesis and PKC activation it is a potentially important target for metabolic drug research. There are three named isoforms.

**[Inhibitor](#) · [Natural Compounds](#) · [Compound Libraries](#) · [Recombinant Proteins](#)**

**This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use**

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481