

Anti-Phospho-PEA15 (Ser116) Polyclonal Antibody 2

Product Details

Ig Type:	IgG
Reactivity:	Rat (predicted:Human,Mouse,Dog,Cow,Rabbit)
Molecular Weight:	Theoretical: 15 kDa.
Purification:	Protein A purified

Applications

Verified Activity:	1. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-Phospho-PEA15 (Ser116) Polyclonal Antibody, Unconjugated (TMAB-11132) 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining
	2. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-Phospho-PEA15 (Ser116) Polyclonal Antibody, Unconjugated (TMAB-11132) 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining
Application:	IHC-P,IHC-Fr,IF
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated Synthesised phosphopeptide: human PEA15 around the phosphorylation site of Ser116
Antigen Species:	Human
Gene ID:	8682
Uniprot ID:	Q15121

Research Background

PED/PEA 15 (Phosphoprotein Enriched in Diabetes/Phosphoprotein Enriched in Astrocytes 15 kDa) is a widely expressed 15 kDa protein comprised of an N terminal region containing a canonical Death Effector Domain (DED) sequence and a nuclear export signal, and a C terminal region containing two serine phosphorylation sites. PED/PEA 15 has been implicated in the regulation of multiple cellular processes including apoptosis, integrin activation, and insulin sensitive glucose transport in insulin responsive cells. Phosphorylation of both serine 104 (a Protein Kinase C site) and serine 116 (a substrate of CaMKII and Akt) is required for PED/PEA 15 function.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481