

Anti-Phospho-PIM1 (Tyr309) Polyclonal Antibody 2

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat (predicted:Dog,Pig,Cow,Rabbit,Sheep)
Molecular Weight:	Theoretical: 45 kDa.
Purification:	Protein A purified

Applications

Verified Activity:	<p>1. Paraformaldehyde-fixed, paraffin embedded (Rat breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-PIM1) Polyclonal Antibody, Unconjugated (TMAB-11141) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.</p> <p>2. Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-PIM1) Polyclonal Antibody, Unconjugated (TMAB-11141) at 1:500 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining.</p> <p>3. Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (Phospho-PIM1 (Tyr309)) Polyclonal Antibody, Unconjugated (TMAB-11141) at 1: 200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.</p>
Application:	IHC-P,IHC-Fr,IF
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated Synthesised phosphopeptide: human PIM1 around the phosphorylation site of Tyr309
Antigen Species:	Human
Gene ID:	5292
Uniprot ID:	P11309

Research Background

PIM1 is a member of the serine/threonine kinase PIM oncogene family. PIM1 has been implicated in lymphomagenesis, cell proliferation, apoptosis, differentiation and tumourigenesis. The PIM1 protein kinase is upregulated in prostate cancer.

The Pim family serine/threonine protein kinases were first identified in studies examining genes targeted for proviral insertion in murine leukemia virus-induced T lymphomas. Increased levels of Pim kinases predispose cells to lymphomagenesis and enhance the activity of mitogenic proteins such as p100, c-Myb, and Cdc25A. In addition, Pim kinases are also involved in modulation of synaptic strength in neurons and anti-apoptotic signaling in hematopoietic progenitor cells. Pim-3, a member of the proto-oncogene Pim family that expresses serine/threonine kinase activity, shares significant homology with Pim-1 serine/threonine protein kinases. Pim-3 may function as a mediator of synaptic plasticity in the brain and is presumably involved in the anti-apoptosis process and cell cycle progression as well as the proliferation of human hepatoma cell lines. The Pim-3 protein is widely expressed, however no expression is observed in the colon, thymus, or small intestine.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481