

Anti-PRKAG2 Polyclonal Antibody

Product Details

| | |
|-------------------|---|
| Ig Type: | IgG |
| Reactivity: | Rat (predicted:Human,Mouse,Chicken,Dog,Pig,Cow,Horse) |
| Molecular Weight: | Theoretical: 63 kDa. |
| Purification: | Protein A purified |

Applications

| | |
|--------------------|--|
| Verified Activity: | <p>1. Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-PRKAG2/AMPKγ2 Polyclonal Antibody, Unconjugated (TMAB-11736) 1: 200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining</p> <p>2. Paraformaldehyde-fixed, paraffin embedded (rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (PRKAG2) Polyclonal Antibody, Unconjugated (TMAB-11736) at 1: 200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) instructions and DAB staining.</p> |
| Application: | IHC-P,IHC-Fr,IF |
| Recommended | IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500 |

Properties

| | |
|----------------------|---|
| Stability & Storage: | Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles. |
| Shipping: | Shipping with blue ice. |

Antigen Details

| | |
|------------------|--|
| Immunogen: | KLH conjugated synthetic peptide: human PRKAG2 |
| Antigen Species: | Human |
| Gene ID: | 51422 |
| Uniprot ID: | Q9UGJ0 |

Research Background

AMPK is a heterotrimeric complex comprising a catalytic α subunit and regulatory β and γ subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate in vivo hydroxy-methylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK α 1 and AMPK α 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK β 1 encodes a 271 amino acid protein and human AMPK β 2 encodes a 272 amino acid protein. The human AMPK γ 1 gene encodes a 331 amino acid

A DRUG SCREENING EXPERT

protein. Human AMPKg2 and AMPKg3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481