

Anti-PTEN Antibody (6C882)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 44 kDa. Actual: 53 kDa.
Clone:	6C882
Purification:	Protein A purified

Applications

Verified Activity:

1. 25 ug total protein per lane of various lysates (see on figure) probed with PTEN monoclonal antibody, unconjugated (TMAB-11893) at 1: 1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
2. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Mouse Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Rat Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Cervical Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Rat Adrenal Gland; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
10. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by

boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Prostate Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

12. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

13. Paraformaldehyde-fixed, paraffin embedded Human Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PTEN Monoclonal Antibody, Unconjugated (TMAB-11893) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

Application: ELISA,IF,IHC-Fr,IHC-P,WB

Recommended ELISA=1:5000-10000; IF=1:200-500; IHC-Fr=1:200-500; IHC-P=1:200-500; WB=1:1000-2000

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human PTEN

Antigen Species: Human

Gene ID: 5728

Uniprot ID: P60484

Research Background

Potential tumor suppressor. Acts as a phosphoinositide3-phosphatase by regulating PtdIns (3,4,5)P3 levels. Involved in regulation of the AKT1 signaling pathway. The unphosphorylated form cooperates with AIP1 to suppress AKT1 activation. The PTEN/MMAC1 discovers the first to have the suppress of the phosphoric acid enzyme activity cancer gene currently. The gene of PTEN locates the chromosome10q23 area, sending forth sex tumor and a few households cancers with the variety to suffer from the comprehensive disease easily relevant. The activity that passes to repress the Akt regulates the cell period, the cell ground rule decrease and glues to connect. This text discussed PTEN structure, function and its correlationses, the PTEN is in tumor repress function mechanism.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481