

Anti-PTGER2 Antibody (10173)

Product Details

Ig Type:	IgG/Kappa
Reactivity:	Human,Mouse,Rat
Clone:	10173
Purification:	Protein A purified

Applications

- 1. Blocking buffer: 5% NFDM/TBST**
Primary ab dilution: 1:1000
Primary ab incubation condition: 4°C overnight
Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)
Lysate: HeLa, JAR, F9, Rat liver
Protein loading quantity: 20 µg
Exposure time: 10 s
Predicted MW: 52 kDa
Observed MW: 56 kDa
- 2. Tissue: Rat lung**
Section type: Formalin fixed & Paraffin-embedded section
Retrieval method: High temperature and high pressure
Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100
Primary ab incubation condition: 1 hour at room temperature
Secondary ab: SP Kit (Rabbit)
Counter stain: Hematoxylin (Blue)
Comment: Color brown is the positive signal for TMAB-11897
- 3. Tissue: Human placenta**
Section type: Formalin fixed & Paraffin-embedded section
Retrieval method: High temperature and high pressure
Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100
Primary ab incubation condition: 1 hour at room temperature
Secondary ab: SP Kit (Rabbit)
Counter stain: Hematoxylin (Blue)
Comment: Color brown is the positive signal for TMAB-11897
- 4. Tissue: Mouse liver**
Section type: Formalin fixed & Paraffin-embedded section
Retrieval method: High temperature and high pressure
Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100
Primary ab incubation condition: 1 hour at room temperature
Secondary ab: SP Kit (Rabbit)
Counter stain: Hematoxylin (Blue)
Comment: Color brown is the positive signal for TMAB-11897
- 5. Cell line: HeLa**
Fixative: 100% Ice-cold methanol
Permeabilization: 0.1% TritonX-100
Primary ab dilution: 1:50
Primary incubation condition: 4°C overnight

Verified Activity:

A DRUG SCREENING EXPERT

Secondary ab: Goat Anti-Rabbit IgG
Nuclear counter stain: DAPI (Blue)
Comment: Color green is the positive signal for TMAB-11897

Application: WB,IHC-P,IHC-Fr,ICC/IF,IF

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:50-200; IF: 1:100-500

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Gene ID: 5732

Research Background

Prostaglandins are produced by the metabolism of arachidonic acid. Prostaglandin E2 is one of the five physiologically significant prostanoids known. Its wide spectrum of physiologic and pharmacologic effects in various tissues is mediated through binding to the Prostaglandin E2 receptors (EP1, EP2, EP3 & EP4). These include effects on the immune, endocrine, cardiovascular, renal and reproductive systems as well as smooth muscle. It is also one of the most abundant of the prostanoid family in the brain where it plays an important role in many neural functions, particularly in newborn babies, and as a mediator of inflammation. Prostaglandin E2 signals through a family of G-protein coupled receptors known as EP receptors. There are 4 subtypes of EP receptors, known as EP1, EP2, EP3 and EP4. EP2 receptors are 358 amino acid proteins with a short third intracellular loop. EP2 receptors stimulate adenylyl cyclase by their coupling to Gs and do not undergo Prostaglandin E2 induced internalization. The EP2 receptors is involved with the contraction and relaxation of smooth muscle tissue. These receptors are mainly localized in lung and placental tissues and in smooth muscle.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481
