

Anti-RALBP1 Antibody (81172)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 76 kDa. Actual: 98 kDa.
Clone:	81172
Purification:	Protein A purified

Applications

Verified Activity:

1. 25 ug total protein per lane of various lysates (see on figure) probed with RALBP1 monoclonal antibody, unconjugated (TMAB-12062) at 1: 1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
2. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Mouse Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Rat Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Mouse Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Rat Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
10. Paraformaldehyde-fixed, paraffin embedded Human Stomach; Antigen retrieval by boiling

A DRUG SCREENING EXPERT

in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RALBP1 Monoclonal Antibody, Unconjugated (TMAB-12062) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

Application: FCM,IF,IHC-Fr,IHC-P,WB

Recommended FCM=1:50-100; IF=1:100-500; IHC-Fr=1:100-500; IHC-P=1:100-500; WB=1:500-2000

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human RalBP1

Antigen Species: Human

Gene ID: 10928

Uniprot ID: Q15311

Research Background

Multifunctional protein that functions as a downstream effector of RALA and RALB. As a GTPase-activating protein/GAP can inactivate CDC42 and RAC1 by stimulating their GTPase activity. As part of the Ral signaling pathway, may also regulate ligand-dependent EGF and insulin receptors-mediated endocytosis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481
