

## Anti-RGS17 Polyclonal Antibody

## Product Details

Ig Type:	IgG
Reactivity:	Rat (predicted: Human, Mouse, Dog, Pig, Cow, Horse, Rabbit, Guinea Pig)
Molecular Weight:	Theoretical: 24 kDa.
Purification:	Protein A purified

## Applications

Verified Activity:	<p>1. Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01 M, pH 6.0), Boiling bathing for 15 min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30 min; Blocking buffer (normal goat serum) at 37°C for 20 min; Incubation: Anti-RGS17 Polyclonal Antibody, Unconjugated (TMAB-12225) 1:400, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining</p> <p>2. Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (RGS17) Polyclonal Antibody, Unconjugated (TMAB-12225) at 1:400 overnight at 4°C, followed by a conjugated secondary for 20 minutes and DAB staining</p>
Application:	IHC-P, IHC-Fr, IF
Recommended	IHC-P: 1:100-500; IHC-Fr: 1:100-500; IF: 1:100-500

## Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

## Antigen Details

Immunogen:	KLH conjugated synthetic peptide: human RGS17
Antigen Species:	Human
Gene ID:	26575
Uniprot ID:	Q9UGC6

## Research Background

This gene encodes a member of the regulator of G-protein signaling family. This protein contains a conserved, 120 amino acid motif called the RGS domain and a cysteine-rich region. The protein attenuates the signaling activity of G-proteins by binding to activated, GTP-bound G alpha subunits and acting as a GTPase activating protein (GAP), increasing the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. [provided by RefSeq, Jul 2008]

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481